

Pressure Damage

Information leaflet
for patients/carers



Advice and Support

We hope that this leaflet will make you aware of how to recognise early signs of pressure damage and the steps you can take to prevent it from occurring.

If you feel you need more information or have any concerns please contact your GP or community nurses.

Contact details for your community nurse are:

Get the best from your NHS

Our patient advice and liaison service (PALS) can help patients, families and carers with questions, comments or concerns about NHS services. If you need advice about our services, facilities or staff, or would like to make a comment, please contact PALS at:



PALS Sussex Community NHS
Foundation Trust Freepost (BR117),
Elm Grove, Brighton BN2 3EW



01273 242292



sc-tr.serviceexperience@nhs.net

Please ask any member of the team caring for you if you need this information in large print, Braille, easy read, audio tape, email. Or if you need help with understanding this information or require this in a language that is not English.



*Excellent care at the
heart of the community*



What is pressure damage?

Pressure damage is a form of damage to the skin and underlying tissue. It is also known as 'pressure ulcers', 'pressure sores', or 'bed sores'.

If untreated it can get worse and seriously affect a person's health and delay their recovery.

Fortunately, most pressure damage can be prevented.

What causes pressure damage?

Pressure damage is caused by a combination of:

Pressure: normal body weight can squash the skin and damage the blood supply to the area. Lying in one position for a long time can cause this.


Slipping/ sliding down a bed or chair: can damage the skin and deeper layers of tissue.

Moisture: urine, faeces and sweat can irritate the skin making it breakdown more easily.

What parts of the body can get pressure damage?

Pressure Damage can develop anywhere on the body but are mostly found over bony areas such as heels, elbows, hips, bottoms, spine and ankles.

They may also occur from items pressing down on the skin such as, tubing from equipment, glasses, or material folds from clothing.



Who is most at risk of developing pressure damage?

You may be at risk of developing pressure damage for a number of reasons including:

- Problems with movement
- Poor circulation
- Some conditions and some treatments may reduce your sensitivity to pain or discomfort which means you are less likely to move or notice pressure damage developing.
- Inadequate diet or fluid intake
- Older skin being thinner and slower to heal.

People with these issues who then become suddenly unwell are extremely vulnerable of pressure damage development.

Early signs of pressure damage

- Pain or discomfort in the skin of any area under pressure, or over a bony area.
- Change in skin colour. Skin may develop reddened areas. Patients with darkly pigmented skin may develop a purple/blue patch.
- Hot or cold patches
- Blistering

Without treatment the damage may get worse, developing into hard, black skin or becoming an open wound.

How can you prevent pressure damage?

Your healthcare team will assess your general health which will help them establish if you are at risk of pressure damage. If necessary, the relevant preventative care will be planned.

However, you can also do the following things to help prevent pressure damage:

Move

- Change your position regularly.
- Use the flat of your foot not your heel when pushing yourself up the bed.
- If you have difficulty moving, ask for help.

Look

Check your skin for pressure damage at least once a day.

Report

Tell a nurse or carer if you are experiencing pain or discomfort in any area of skin.

Protect

Wash your skin every day using a mild soap and warm water.

Inform

Tell your health care team if you suffer from incontinence as they can provide pads to prevent soreness.

Eat a well-balanced diet

Make sure you eat a balanced diet and drink plenty of fluids.